


Journal Pre-proof

Diez años de experiencia en el abordaje conjunto,
dermatología-reumatología, de los pacientes con artritis psoriásica

A. López Ferrer H. Corominas Macias L. Puig Sanz A. Laiz Alonso



PII: S0001-7310(25)00457-0

DOI: <https://doi.org/doi:10.1016/j.ad.2025.06.013>

Reference: AD 4414

To appear in: *Actas dermosifiliograficas*

Received Date: 22 July 2024

Accepted Date: 6 January 2025

Please cite this article as: López Ferrer A, Corominas Macias H, Puig Sanz L, Laiz Alonso A, Diez años de experiencia en el abordaje conjunto, dermatología-reumatología, de los pacientes con artritis psoriásica, *Actas dermosifiliograficas* (2025), doi: <https://doi.org/10.1016/j.ad.2025.06.013>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 AEDV. Published by Elsevier España, S.L.U.

Refers to AD_4343

Carta científico-clínica

Diez años de experiencia en el abordaje conjunto, dermatología-reumatología, de los pacientes con artritis psoriásica

[[Translated article]] Ten-Year Experience in the Joint Dermatology-Rheumatology Approach to Patients With Psoriatic Arthritis

A. López Ferrer^a, H. Corominas Macias^b, L. Puig Sanz^c, A. Laiz Alonso^d.

- a. Servicio de Dermatología, Hospital de la Santa Creu i Sant Pau, Barcelona. Barcelona.
- b. Servicio de Reumatología, Hospital de la Santa Creu i Sant Pau, Barcelona. Barcelona.
- c. Servicio de Dermatología, Hospital de la Santa Creu i Sant Pau, Barcelona. Barcelona.
- d. Servicio de Reumatología, Hospital de la Santa Creu i Sant Pau, Barcelona. Barcelona.

Autor de correspondencia:

Anna López Ferrer

alopezfe@santpau.cat

To the Editor,

Joint management of psoriatic arthritis (PsA) can be very effective in some patients, although there is limited evidence in the literature based on large series of clinical data¹⁻⁹.

The aim of this article is to quantify the diagnostic and therapeutic changes experienced by patients due to their passage through the PAIDER clinic (Comprehensive Care Program for Dermatology and Rheumatology Patients) of a tertiary referral center specializing in the management of immunomediated diseases from June 29th, 2012 through July 4th, 2023. This study was approved by the center ethics committee. An episode was defined as the set of visits made by a patient until discharge. A complex patient was defined as one who exhibited psychiatric, cardiac, hepatic, or oncological disease, communication problems, or treatment-related issues. Statistical analysis was performed using SPSS 28 statistical software.

The sample consists of 970 episodes generated from 732 patients who attended the clinic on different occasions over the years. A total of 53.8% of these were women, and the mean age was 53.1 years. Disease duration, origin, reasons for consultation detailed by referring services, comorbidities, and the proportion of complex patients are shown in Tables 1 and 2.

The median number of visits per episode was 1, with a minimum of 0 (patient did not attend) and a maximum of 22 visits, interquartile range (IQR) = (1.0; 2.0).

Diagnostic changes showed an increase in favor of PsA. Among the patients' initial diagnoses, we found 381 (40.4%) with some type of PsA (axial, peripheral, or mixed) and 378 (40%) with psoriasis. Other diagnoses included rheumatoid arthritis (RA), ankylosing spondylitis (AS), SAPHO (synovitis, acne, pustulosis, hyperostosis, and osteitis), and hidradenitis suppurativa (HS). Patients with diagnoses of dactylitis, enthesitis, and/or mono-oligoarthritis were grouped under the same heading. Final diagnoses showed that 613 (65.2%) patients had some type of PsA (axial, peripheral, or mixed), 217 (23.1%) had psoriasis, and the rest showed other diagnoses (Table 1).

In 329 episodes of patients referred for dermatological evaluation, the diagnoses were psoriasis in 154 (46.8%), psoriatic onychopathy in 35 (10.6%), onychomycosis in 18 (5.5%), and others in 122 (37.1%). Of these, 187 (56.8%) served to confirm the suspicion of PsA. In 432 episodes of patients referred for rheumatological evaluation, the diagnoses were arthritis in 161 (37.3%), enthesitis in 39 (9%), dactylitis in 18 (4.2%), osteoarthritis in 93 (21.5%), crystalline arthritis in 20 (23.4%), and others in 101 (37.1%). Of these, 201 (46.5%) served to confirm the suspicion of PsA (Table 3).

Patients could present one or more diagnoses from the dermatologist or rheumatologist, in the same episode or successive episodes.

Changes in patient treatment occurred in 430 episodes (46.3%): treatment was initiated in 137 (14.7%), an additional treatment was added to the existing one in 102 (11%), a treatment change within the same therapeutic class was made in 93 (10%), a change of therapeutic class was made in 67 (7.2%), and treatment was stopped in 31 (3%).

As far as we know, this is the largest cohort of patients under joint follow-up in a dermatology-rheumatology clinic. The work describes that more than half experience a change in diagnosis and/or treatment. In our case, the number of new PsA cases increases among patients with an initial diagnosis of psoriasis and, to a lesser extent, with other musculoskeletal symptoms (oligoarthritis, dactylitis, enthesitis...), likely because most of our patients first developed skin signs, as is common in the literature¹⁰.

Regarding treatment changes, our data show that these were performed in patients from rheumatology, indicating that the reason was mostly skin involvement. In addition, the approval processes for some drugs can take years of difference between specialties, often favoring dermatology, which means that patients referred to these clinics may have a wider therapeutic arsenal available.

Limitations of this study include the fact that it is a single-center cohort, where patients from smaller hospitals may have been referred due to their complexity, which could lead to referral bias. However, this study includes a large sample size of consecutive patients, referred to a joint clinic and with prolonged follow-up. These results show that multidisciplinary care is associated with a change in diagnosis, fundamentally in favor of PsA and, linked to this, a change in treatment, with the aim of improving the prognosis of the disease.

Funding

This work received funding from UCB Pharma for the statistical analysis of the data.

Conflicts of interest

A. López Ferrer has received payments or honoraria for educational events, support to attend meetings, and participation in advisory boards or consultancies from Abbvie, Almirall, Amgen, Boehringer Ingelheim, Bristol Myers Squibb Janssen, Eli Lilly, Leo Pharma, Novartis, UCB.

H. Corominas has received payments or honoraria for educational events, support to attend meetings, and participation in advisory boards or consultancies from Grünenthal, MSD, Biogen, Galapagos, Abbvie, Roche, Bristol Myers Squibb, Gebro, Sanofi, and UCB.

L. Puig Sanz has received payments or honoraria for educational events, support to attend meetings, and participation in advisory boards or consultancies from Abbvie, Almirall, Amgen, Biogen, Boehringer Ingelheim, Bristol Myers Squibb, Dice, Janssen, Leo-Pharma, Lilly, Novartis, Pfizer, Sandoz, Sanofi, and UCB.

A. Laiz Alonso has received payments or honoraria for educational events, support to attend meetings, and participation in advisory boards or consultancies from Abbvie, Almirall, Amgen, Janssen, MSD, Novartis, Pfizer, and UCB.

Acknowledgments

The authors wish to thank the members of the Dermatology and Rheumatology services of *Hospital de la Santa Creu and Sant Pau* for their participation in the study and for referring patients to the joint PAIDER clinic, and especially Dr. Eva Vilarrasa, Dr. Hye Park, and Dr. Patricia Moya for their close collaboration with the joint management of patients in our clinic.

References

<BIBL>

<BIB>

1

S Okhovat F J.P.

S Ogdie F A.

S Reddy F S.M.

S Rosen F C.F.

S Scher F J.U.

S Merola F J.F.

AT Psoriasis and Psoriatic Arthritis Clinics Multicenter Advancement Network Consortium (PPACMAN) Survey: Benefits and Challenges of Combined Rheumatology-dermatology Clinics

JT J Rheumatol.

V 44

D 2017

P 693-L 694

DOI 10.3899/jrheum.170148

<original_ref>[1] Okhovat JP, Ogdie A, Reddy SM, Rosen CF, Scher JU, Merola JF. Psoriasis and Psoriatic Arthritis Clinics Multicenter Advancement Network Consortium (PPACMAN) Survey: Benefits and Challenges of Combined Rheumatology-dermatology Clinics. J Rheumatol. 2017 May 1;44<I>(5):693–4.

<DOI>10.3899/jrheum.170148.</DOI> </original_ref>

</BIB>

<BIB>

2

S Urruticoechea-Arana F A.

S Serra Torres F M.

S Hergueta Diaz F M.

S González Guerrero F M.E.

S Fariñas Padron F L.

S Navarro Martín F S.<ET-AL>

AT Experience and satisfaction with a multidisciplinary care unit for patients with psoriasis an psoriatic arthritis

JT Reumatol Clin.

V 15

D 2019

P 237-L 241

DOI 10.1016/j.reuma.2017.07.007

<original_ref>[2] Urruticoechea-Arana A, Serra Torres M, Hergueta Diaz M, González Guerrero ME, Fariñas Padron L, Navarro Martín S, et al. Experience and satisfaction with a multidisciplinary care unit for patients with psoriasis an psoriatic arthritis. Reumatol Clin. 2019;15<I>(4):237–41.

<DOI>10.1016/j.reuma.2017.07.007.</DOI> </original_ref>

</BIB>

<BIB>

3

S Carubbi F F.

S Chimenti F M.S.

S Blasetti F G.

S Cipriani F P.

S Musto F A.
 S Fagnoli F M.C.<ET-AL>
 AT Association of psoriasis and/or psoriatic arthritis with autoimmune diseases: the experience of two Italian integrated Dermatology/Rheumatology outpatient clinics
 JT J Eur Acad Dermatol Venereol.
 V 29
 D 2015
 P 2160-L 2168
 DOI 10.1111/jdv.13170
 <original_ref>[3] Carubbi F, Chimenti MS, Blasetti G, Cipriani P, Musto A, Fagnoli MC, et al. Association of psoriasis and/or psoriatic arthritis with autoimmune diseases: the experience of two Italian integrated Dermatology/Rheumatology outpatient clinics. J Eur Acad Dermatol Venereol. 2015;29<I>(11):2160–8.
 <DOI>10.1111/jdv.13170.</DOI></original_ref>
 </BIB>

<BIB>
 4
 S Queiro F R.
 S Coto F P.
 S Joven F B.
 S Rivera F R.
 S Navío Marco F T.
 S de la Cueva F P.<ET-AL>
 AT Current status of multidisciplinary care in psoriatic arthritis in Spain: NEXUS 2.0 project
 JT Reumatol Clin.
 V 16
 D 2020
 P 24-L 31
 <original_ref>[4] Queiro R, Coto P, Joven B, Rivera R, Navío Marco T, de la Cueva P, et al. Current status of multidisciplinary care in psoriatic arthritis in Spain: NEXUS 2.0 project. Reumatol Clin. 2020;16<I>(1):24–31.</original_ref>
 5

S Cañete F J.D.
 S Puig F L.
 AT Multidisciplinary teams for psoriatic arthritis: on aims and approaches
 JT Actas Dermosifiliogr.
 V 105
 D 2014
 P 325-L 327
 <original_ref>[5] Cañete JD, Puig L. Multidisciplinary teams for psoriatic arthritis: on aims and approaches. Actas Dermosifiliogr. 2014 May;105<I>(4):325–
 7.</original_ref>
 </BIB>

<BIB>
 6
 S Gratacós F J.
 S Behrens F F.
 S Coates F L.C.
 S Lubrano F E.
 S Thaçi F D.
 S Bundy F C.<ET-AL>

AT A 12-point recommendation framework to support advancement of the multidisciplinary care of psoriatic arthritis: A call to action

JT Joint Bone Spine.

V 88

D 2021

P 105175

<original_ref>[6] Gratacós J, Behrens F, Coates LC, Lubrano E, Thaçi D, Bundy C, et al. A 12-point recommendation framework to support advancement of the multidisciplinary care of psoriatic arthritis: A call to action. Joint Bone Spine. 2021;88<I>(3):105175.</original_ref>

7

S Luelmo F J.

S Gratacós F J.

S Moreno Martínez-Losa F M.

S Ribera F M.

S Romaní F J.

S Calvet F J.<ET-AL>

AT A report of 4 years of experience of a multidisciplinary unit of psoriasis and psoriatic arthritis

JT Reumatol Clin.

V 10

D 2014

P 141-L 146

DOI 10.1016/j.reuma.2014.01.004

<original_ref>[7] Luelmo J, Gratacós J, Moreno Martínez-Losa M, Ribera M, Romaní J, Calvet J, et al. A report of 4 years of experience of a multidisciplinary unit of psoriasis and psoriatic arthritis. Reumatol Clin. 2014;10<I>(3):141–6.

<DOI>10.1016/j.reuma.2014.01.004.</DOI></original_ref>

</BIB>

<BIB>

8

S Velez F N.F.

S Wei-Passanese F E.X.

S Husni F M.E.

S Mody F E.A.

S Qureshi F A.A.

AT Management of psoriasis and psoriatic arthritis in a combined dermatology and rheumatology clinic

JT Arch Dermatol Res [Internet].

V 304

D 2012

P 7-L 13

DOI 10.1007/s00403-011-1172-6

<original_ref>[8] Velez NF, Wei-Passanese EX, Husni ME, Mody EA, Qureshi AA. Management of psoriasis and psoriatic arthritis in a combined dermatology and rheumatology clinic. Arch Dermatol Res [Internet]. 2012;304<I>(1):7–13.

<DOI>10.1007/s00403-011-1172-6.</DOI></original_ref>

</BIB>

<BIB>

9

S Von Ahnen F J.A.

S Gross F G.

S Lull F C.

S Blauth F M.

S Kraemer F B.

S Olsavszky F V.<ET-AL>

AT The Effect of an Interdisciplinary Dermatological-Rheumatological Consultation on the Outcome of Patients with Psoriasis with Musculoskeletal Pain: A Prospective, Single-Center Cohort Study

JT Adv Ther.

V 40

D 2023

P 5243-L 5253

DOI 10.1007/s12325-023-02667-y

<original_ref>[9] von Ahnen JA, Gross G, Lull C, Blauth M, Kraemer B, Olsavszky V, et al. The Effect of an Interdisciplinary Dermatological-Rheumatological Consultation on the Outcome of Patients with Psoriasis with Musculoskeletal Pain: A Prospective, Single-Center Cohort Study. Adv Ther. 2023;40<I>(12):5243–53.

<DOI>10.1007/s12325-023-02667-y.</DOI></original_ref>

</BIB>

<BIB>

10

S Karmacharya F P.

S Wright F K.

S Achenbach F S.J.

S Crowson F C.S.

S Ogdie F A.

S Bekele F D.<ET-AL>

AT Time to transition from psoriasis to psoriatic arthritis: A population-based study

JT Semin Arthritis Rheum.

V 1

D 2022

P 52

DOI 10.1016/j.semarthrit.2021.12.013

<original_ref>[10] Karmacharya P, Wright K, Achenbach SJ, Crowson CS, Ogdie A, Bekele D, et al. Time to transition from psoriasis to psoriatic arthritis: A population-based study. Semin Arthritis Rheum. 2022 Feb 1;52.

<DOI>10.1016/j.semarthrit.2021.12.013.</DOI></original_ref>

Table 1. Clinical and Demographic Characteristics of Patients and Initial and Final Diagnoses in the PAIDER Clinic

Clinical and Demographic Characteristics (n = 970)

<i>Age (mean, years)</i>	53.1	
<i>Female, n (%)</i>	522	53.8
<i>Male, n (%)</i>	448	46.2
<i>Mean disease duration - Dermatology (years)</i>	16.8	
<i>Mean disease duration - Rheumatology (years)</i>	7.5	
<i>Mean number of visits</i>	2.2	
<i>N (%)</i>		
<i>Referring Department^a</i>		
Primary Care	54	5.6
Dermatology	437	45.1
Rheumatology	434	44.7
Other specialties	45	4.6
<i>Reason for Referral</i>	n=968	
Diagnosis	430	44.4
Treatment	253	26.1
Both	285	29.4
<i>Comorbidities</i>	n=959	53.1
Dyslipidemia	280	29.2
Diabetes Mellitus	124	12.9
Hypertension	268	28
Hyperuricemia	105	11.1
Obesity (BMI > 30)	309	33.7
Smoking	249	26.3
Inflammatory Bowel Disease	39	18.9
Osteoporosis	99	18.7
Uveitis	32	3.5
<i>Complexity</i>	n=963	54.3
Cardiac Dysfunction	75	7.8
Hepatic Dysfunction	178	18.5
Mental Disorder	203	21.1
Neoplasm	78	8.1
Adverse Effects	151	15.7
Paradoxical Reaction	89	9.2
Communication Problem	60	6.2
<i>Diagnosis, n (%)</i>	Initial	Final
Peripheral PsA	243 (25.7)	413 (43.9)
Axial PsA	45 (4.8)	79 (8.4)
Mixed PsA	93 (9.9)	121 (12.9)
Rheumatoid Arthritis	24 (2.5)	27 (2.9)
Ankylosing Spondylitis	48 (5.1)	24 (2.5)
SAPHO	11 (1.2)	12 (1.3)
Hidradenitis Suppurativa	73 (0.3)	7 (0.7)
Dactylitis/Enthesitis/Mono-Oligoarticular Arthritis	99 (10.5)	41 (4.3)
Psoriasis	378 (40)	217 (23.1)

^a Most referrals for diagnosis came from dermatology and rheumatology, while referrals for both reasons predominantly came from primary care and other specialties. Differences were statistically significant ($p < 0.001$).

Table 2. Reasons for Consultation by Referring Service

	Consultation for Diagnosis, n (%)	Consultation for Treatment, n (%)	Consultation for Diagnosis and Treatment, n (%)
Primary Care	14 (25.9)	11 (20.4)	29 (53.7)
Dermatology	220 (50.3)	79 (18.1)	138 (31.6)
Rheumatology	191 (44.0)	152 (35.0)	91 (21.0)
Other Hospital Specialists	5 (11.6)	11 (25.5)	27 (62.7)

Data from 968 episodes.

Table 3. Dermatological and Rheumatological Diagnoses in the PAIDER Clinic

Dermatological Disease in Patients with Arthritis n = 329, n (%)

Psoriasis	154 (46.8)
Nail Psoriasis	35 (10.6)
Onychomycosis	18 (5.5)
Others	122 (37.1)

Rheumatological Disease in Patients with Psoriasis n = 432, n (%)

Arthritis	161 (37.3)
Enthesitis	39 (9)
Dactylitis	18 (4.2)
Osteoarthritis	93 (21.5)
Gout	20 (23.4)
Others	101 (37.1)