

Journal Pre-proof

Massive facial involvement in crusted (Norwegian) scabies

Alessandra Michelucci MD Cristian Fidanzi MD Valentina Dini MD
PhD



PII: S0001-7310(25)00471-5

DOI: <https://doi.org/doi:10.1016/j.ad.2024.02.049>

Reference: AD 4428

To appear in: *Actas dermosifiliograficas*

Received Date: 16 January 2024

Accepted Date: 6 February 2024

Please cite this article as: Michelucci A, Fidanzi C, Dini V, Massive facial involvement in crusted (Norwegian) scabies, *Actas dermosifiliograficas* (2025), doi: <https://doi.org/10.1016/j.ad.2024.02.049>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier España, S.L.U. on behalf of AEDV.

Sección: Imágenes en Dermatología

English title: **Massive facial involvement in crusted (Norwegian) scabies**

título en español: **Afectación facial masiva en la sarna costrosa (noruega)**

Alessandra Michelucci MD¹, Cristian Fidanzi MD^{1,2}, and Valentina Dini MD, PhD¹

¹ Unit of Dermatology, University of Pisa, Pisa, Italy

² Unit of Dermatology, Hospital of Carrara, Carrara, Italy

Corresponding author:

Cristian Fidanzi, MD

E-mail address: cri.fidanzi@outlook.it -

An 82-year-old woman presented with hyperkeratotic erythrodermia and onyxia, involving the face (Fig. A), scalp, and palmoplantar region. She reported no comorbidities or personal history of other skin diseases. The lesions were completely asymptomatic, and the patient did not report any painful or itchy symptoms. The lesions had been present for approximately 3 months, progressively worsening despite therapy with emollients initially and later with topical and oral corticosteroids (oral methylprednisolone 16 mg).

The visualization of characteristic burrows during dermoscopic examination of the lesions (Fig. B, mites indicated by arrows) allowed us to establish a diagnosis of crusted scabies (Norwegian scabies). Dermoscopic evaluation should be conducted in regions with less hyperkeratosis, and the identification of the "jet in the sky" sign at the subungual or interdigital region is crucial for diagnosis.

Among the various *Sarcoptes scabiei* variants, crusted (Norwegian) scabies stands out as a highly contagious form, especially affecting immunocompromised individuals, which promotes widespread parasite proliferation. The occasional complete absence of itching and the extensive involvement of the face and scalp, distinguishing it from the classic clinical form, can lead to delayed diagnosis. Frequent misdiagnosis with other conditions, such as eczema, psoriasis rupioides, seborrheic dermatitis, or fungal infections, can also result in incorrect or even worsening treatments of the disease.

Fig.1

