

Acknowledgments

We wish to thank all patients who generously shared their experience for the purposes of this article in the spirit of helping other women.

References

1. Virgili A, Borghi A, Toni G, Minghetti S, Corazza M. Prospective clinical and epidemiologic study of vulvar lichen sclerosis: Analysis of prevalence and severity of clinical features, together with historical and demographic associations. *Dermatology*. 2014;228:145–51.
2. Lee A, Bradford J, Fischer G. Long-term management of adult vulvar lichen sclerosis. A prospective cohort study of 507 women. *JAMA Dermatol*. 2015;151:1061–7.
3. Belotto RA, Chavantes MC, Tardivo JP, dos Santos RE, Fernandes RCM, Horliana ACRT. Therapeutic comparison between treatments for vulvar lichen sclerosis: Study protocol of a randomized prospective and controlled trial. *BMC Womens Health*. 2017;17:61.
4. Kirtschig G, Becker K, Günthert A, Jasaitiene D, Cooper S, Chi CC. Evidence-based (S3) guideline on (anogenital) lichen sclerosis. *J Eur Acad Dermatol Venereol*. 2015;29:e1–43.
5. Mazdziarz A, Osuch B, Kowalska M, Nalewczynska A, Spiewankiewicz B. Photodynamic therapy in the treatment of vulvar lichen sclerosis. *Photodiagnosis Photodyn Ther*. 2017;19:135–9.
6. Rangatchew F, Knudsen J, Thomsen MV, Drzewiecki KT. Surgical treatment of disabling conditions caused by anogenital lichen sclerosis in women: An account of surgical procedures and

results, including patient satisfaction, benefits, and improvements in health-related quality of life. *J Plast Reconstr Aesthet Surg*. 2017;70:501–8.

7. Gurumurthy M, Morah N, Gioffre G, Cruickshank ME. The surgical management of complications of vulvar lichen sclerosis. *Eur J Obstet Gynecol Reprod Biol*. 2012;162:79–82.
8. Hackenjos K, Schröder W, Schöpf E, Vanscheidt W. Therapy of lichen sclerosis et atrophicus vulvae with the CO₂ silk touch laser. *Hautarzt*. 2000;51:502–4.
9. Lee A, Lim A, Fischer G. Fractional carbon dioxide laser in recalcitrant vulvar lichen sclerosis. *Australas J Dermatol*. 2016;57:39–43.
10. Mendieta-Eckert M, Ocerin-Guerra I, Landa-Gundin N. Lichen sclerosis et atrophicus in a surgical scar treated with fractional laser. *J Cosmet Laser Ther*. 2017;19:106–8.

M. Mendieta-Eckert,^{a,*} J. Torrontegui Bilbao,^b
I. Zabalza Estévez,^c N. Landa Gundin^a

^a Servicio de Dermatología, Clínica Dermitek, Bilbao, Spain

^b Equipo enfermería, Clínica Dermitek, Bilbao, Spain

^c Servicio de Anatomía Patológica, Clínica Dr Guimón, Bilbao, Spain

* Corresponding author.

E-mail address: mmendieta@dermitek.com
(M. Mendieta-Eckert).

<https://doi.org/10.1016/j.adengl.2020.12.001>
1578-2190/ © 2020 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Are Patch Test Results Usually Relevant in Patients With Burning Mouth Syndrome?☆



¿Suelen ser relevantes las pruebas epicutáneas en el paciente con síndrome de boca urente?

To the Editor:

Most publications on the results of patch testing in patients with burning mouth syndrome (BMS) report high percentages of positive “clinically relevant” findings (between 13% and 74%), thus leading this approach to be considered a useful tool in the assessment of affected patients.^{1–3} Some authors point out that patients with intermittent symptoms (type 3 BMS) could be ideal candidates for patch testing.³ Various allergens are involved, and these are not always consistent from study to study, although most positive results are for metals, especially nickel.

Taking the above into account, we performed a retrospective observational study in order to determine the clinical relevance of positive patch test results in patients with BMS. We analyzed the results of patch tests performed in patients diagnosed with BMS referred to our skin allergy unit over a period of 15 years (January 2001 to December 2015). We also followed patients via their electronic health records (Osabide Global) in order to determine the relevance of the positive results. During the period analyzed, a total of 2789 patients underwent testing. Of these, 67 had consulted for BMS (Fig. 1). The characteristics of the cohort were reported using the MOAHLFA index (Table 1). All patients with BMS underwent testing with the standard series of the Spanish Contact Dermatitis and Skin Allergy Research Group (Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea [GEIDAC]). Supplementary series were applied in 65 cases, depending on the clinical history. The most common were a dental series (Chemotechnique) (52.23%) and a metal series (Chemotechnique) (44.7%) (Table 2). Readings were made at 48 and at 96 hours following the recommendations of the European Society of Contact Dermatitis (ESCD). An allergic reaction was observed in 23 of the 67 patients (34.3%), with a total of 32 positive results recorded. Nickel was the main allergen (n = 11), followed by methylisothiazolinone/methylchlorisothiazolinone (n = 3) and beryllium (n = 3). We were able to establish present relevance in

☆ Please cite this article as: De la Torre Gomar FJ, Heras González S, González Pérez R. ¿Suelen ser relevantes las pruebas epicutáneas en el paciente con síndrome de boca urente? *Actas Dermosifiliogr*. 2021;112:88–89.




Figure 1 Patients with burning mouth syndrome and positive patch test results. In 2 of the 67 affected patients (2.9%), we established present relevance for the positive results, both of which were to nickel (Ni).

Table 1 MOAHLFA Index of the Study Cohort.

	n = 2789	BMS
M (Male)	979	13
O (Occupational) ^a	60	0
A (Atopy)	238	0
H (Hand) ^a	714	0
L (Leg) ^a	174	0
F (Face) ^a	356	0
A (Aged > 40)	2144	67

^a The items of the MOAHLFA index corresponding to O, H, L, and F are not applicable with respect to the presenting complaint (burning mouth syndrome [BMS]), given that it is not considered to be occupational and does not affect the hands, legs, or face.

Table 2 Series Used in the Patch Tests Carried Out.

Series	Number of Patients	Percentage of the Sample
Standard	67	100%
Dental	35	52.23%
Metal	30	44.78%
Bakery	16	23.88%
Own products	7	10.45%
Cosmetics	1	1.49%

only 2 of the 67 patients with BMS (2.9%). In both cases, the patients were sensitized to the nickel in their dental prostheses; the symptoms resolved when the prostheses, which contained nickel, were removed. In the 21 remaining patients whose test results were positive, the results were considered to be not relevant or of unknown relevance for various reasons, mainly because of positive results to metals that were not a component of the dental prostheses they were wearing at the time (Fig. 1).

In the present study cohort, only 2.9% of the positive results were considered clinically relevant or with present relevance. This percentage is clearly lower than those

reported in the literature.¹⁻³ The cases considered as having present relevance corresponded to nickel, an allergen of the GEIDAC standard series. Therefore, the supplementary series used were not associated with an increase in the number of clinically relevant positive results diagnosed. Based on these findings, we believe that, except for selected cases where the clinical history points to an allergic origin for symptoms, patch testing should not be included in the routine evaluation of patients with BMS.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

1. Steele JC, Bruce AJ, Davis MD, et al. Clinically relevant patch test results in patients with burning mouth syndrome. *Dermatitis*. 2012;23(2):61–71.
2. Lynde CB, Grushka M, Walsh SR. Burning mouth syndrome: patch test results from a large case series. *J Cutan Med Surg*. 2014;18(3):174–9.
3. Marino R, Capaccio P, Pignataro L, Spadari F. Burning mouth syndrome: the role of contact hypersensitivity. *Oral Dis*. 2009;15(4):255–8.

De la Torre Gomar F.J.,* Heras González S., González Pérez R.

Servicio de Dermatología, Hospital Universitario Araba, Vitoria-Gasteiz, Spain

* Corresponding author.

E-mail address: fjtogo@gmail.com (F.J. De la Torre Gomar).

2 April 2019 13 May 2019

<https://doi.org/10.1016/j.adengl.2019.05.020>
1578-2190/ © 2020 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).